



Tauberian theorems for the (J, p) summability method

İbrahim Çanak^{a,*}, Ümit Totur^b

^a Department of Mathematics, Ege University, Izmir 35100, Turkey

^b Department of Mathematics, Adnan Menderes University, Aydin 09010, Turkey

ARTICLE INFO

Article history:

Received 13 June 2011

Accepted 8 December 2011

Keywords:

Tauberian conditions and theorems

(J, p) summability

Weighted mean summability

Slow oscillation

ABSTRACT

In this paper, we prove Tauberian theorems of slowly oscillating type for the (J, p) summability method by using a result due to Mikhalin [G.A. Mikhalin, Theorem of Tauberian type for (J, p_n) summation methods, Ukrain. Mat. Zh. 29 (1977), 763–770. English translation: Ukrain. Math. J. 29 (1977) 564–569].

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Assume that $p = (p_n)$ be a sequence of nonnegative numbers with $p_0 > 0$, such that

$$P_n := \sum_{k=0}^n p_k \rightarrow \infty \quad \text{as } n \rightarrow \infty,$$

and that

$$p(x) = \sum_{k=0}^{\infty} p_k x^k < \infty \quad \text{for } 0 < x < 1.$$

The weighted means of a sequence (u_n) of real numbers are defined by

$$\sigma_{n,p}^{(1)}(u) := \frac{1}{P_n} \sum_{k=0}^n p_k u_k$$

for all nonnegative integers n .

If

$$\lim_{n \rightarrow \infty} \sigma_{n,p}^{(1)}(u) = s, \tag{1}$$

then we say that (u_n) is summable to s by the weighted mean method determined by $p = (p_n)$ and we write $u_n \rightarrow s (\bar{N}, p)$.

If $\sum_{k=0}^{\infty} p_k u_k x^k$ is convergent for $0 < x < 1$, and $\frac{1}{p(x)} \sum_{k=0}^{\infty} p_k u_k x^k \rightarrow s$ as $x \rightarrow 1^-$, we say that (u_n) is summable to s by the power series method (J, p) , and we write $u_n \rightarrow s (J, p)$.

* Corresponding author. Tel.: +90 232 311 5418; fax: +90 232 388 1036.

E-mail addresses: ibrahimcanak@yahoo.com, ibrahim.canak@ege.edu.tr (İ. Çanak), utotur@adu.edu.tr (Ü. Totur).

The (\bar{N}, p) and (J, p) summability methods are regular if and only if $P_n \rightarrow \infty$ as $n \rightarrow \infty$. It is well known that $u_n \rightarrow s(\bar{N}, p)$ implies $u_n \rightarrow s(J, p)$ (see [1]).

The sequence of the backward differences of $u = (u_n)$ is denoted by $\Delta u = (\Delta u_n)$ and defined by $\Delta u_n = u_n - u_{n-1}$ for $n \geq 1$ and $\Delta u_0 = u_0$. The identity

$$u_n - \sigma_{n,p}^{(1)}(u) = V_{n,p}^{(0)}(\Delta u) \quad (2)$$

where $V_{n,p}^{(0)}(\Delta u) := \frac{1}{P_n} \sum_{k=1}^n P_{k-1} \Delta u_k$. The identity (2) is called the weighted Kronecker identity and $(V_{n,p}^{(0)}(\Delta u))$ is called a generator sequence of (u_n) .

For any nonnegative integer m , we define $\sigma_{n,p}^{(m)}(u)$ and $V_{n,p}^{(m)}(\Delta u)$ by

$$\sigma_{n,p}^{(m)}(u) = \begin{cases} \frac{1}{P_n} \sum_{k=0}^n p_k \sigma_{k,p}^{(m-1)}(u), & m \geq 1 \\ u_n, & m = 0 \end{cases}$$

and

$$V_{n,p}^{(m)}(\Delta u) = \begin{cases} \frac{1}{P_n} \sum_{k=0}^n p_k V_{k,p}^{(m-1)}(\Delta u), & m \geq 1 \\ V_{n,p}^{(0)}(\Delta u), & m = 0 \end{cases}$$

respectively.

If

$$\lim_{n \rightarrow \infty} \sigma_{n,p}^{(m)}(u) = s, \quad (3)$$

then we say that (u_n) is summable to s by the (\bar{N}_m, p) method and we write $u_n \rightarrow s(\bar{N}_m, p)$. If $\sigma_{n,p}^{(m)}(u) \rightarrow s(J, p)$, we write $u_n \rightarrow s(J, p)(\bar{N}_m, p)$ for any nonnegative integer m .

A sequence (u_n) is slowly oscillating [2] if

$$\lim_{\lambda \rightarrow 1^+} \limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta u_j \right| = 0. \quad (4)$$

For the definitions of $O(1)$ and $o(1)$ we refer the reader to [3, page 149].

In this paper our object is to establish the following Tauberian theorems for (J, p) and $(J, p)(\bar{N}_m, p)$ summability methods by using the concept of the generator sequence.

Theorem 1. Let (p_n) satisfy the conditions

$$p_0 > 0, \quad p_n \geq 0 \quad (n = 1, 2, \dots), \quad P_n = \sum_{k=0}^n p_k \rightarrow \infty \quad (n \rightarrow \infty), \quad (5)$$

$$1 \leq \frac{P_n}{n} \rightarrow 1 \quad (n \rightarrow \infty), \quad (6)$$

$$\left(\frac{1}{P_n} \right) \text{ is an absolutely monotone sequence.} \quad (7)$$

Then slow oscillation of (u_n) is a Tauberian condition for (J, p) summability.

Theorem 2. Let (p_n) satisfy the conditions (5)–(7). Then slow oscillation of (u_n) is a Tauberian condition for $(J, p)(\bar{N}_m, p)$ summability.

Note that in Theorems 1 and 2, if we get $p_n = 1$ for all nonnegative integer n , we have the Tauberian theorems for Abel and generalized Abel summability methods obtained in [4].

Our proofs are based on the following result due to Mikhlin [5].

Theorem 3. Let (p_n) satisfy the conditions (5),

$$1 \leq \frac{P_m}{P_n} \rightarrow 1 \quad \text{when } 1 < \frac{m}{n} \rightarrow 1 \quad (n \rightarrow \infty), \quad (8)$$

and (7). If $V_{n,p}^{(0)}(\Delta u) \geq -C$ for some $C \geq 0$, then $u_n \rightarrow s(J, p)$ implies $u_n \rightarrow s(\bar{N}, p)$.

Note that the condition (6) implies the condition (8).

2. Lemmas

We need the following lemmas to prove our theorems.

Lemma 4. If (λ_n) is an increasing sequence of positive real numbers and (v_n) is a sequence of real numbers, then

$$\left| \sum_{k=p}^q \lambda_k \Delta v_k \right| \leq \lambda_q \max_{p \leq r \leq q} \left| \sum_{k=r}^q \Delta v_k \right|.$$

Proof. We have

$$\sum_{k=p}^q \lambda_k (v_k - v_{k-1}) = \sum_{k=p}^q \lambda_k ((v_q - v_{k-1}) - (v_q - v_k)) = \sum_{k=p}^q \lambda_k (S_k - S_{k-1}),$$

where $S_k = \sum_{j=k}^q \Delta v_j$ and $S_{q+1} = 0$.

By summation by parts, we have

$$\begin{aligned} \sum_{k=p}^q \lambda_k (v_k - v_{k-1}) &= \sum_{k=p}^q S_k \lambda_k - \sum_{k=p}^q S_{k+1} \lambda_k \\ &= \sum_{k=p}^q S_k \lambda_k - \sum_{k=p+1}^{q+1} S_k \lambda_{k-1} \\ &= \sum_{k=p+1}^q S_k (\lambda_k - \lambda_{k-1}) + S_p \lambda_p - S_{q+1} \lambda_q. \end{aligned}$$

Since (λ_n) is an increasing sequence of positive real numbers, we obtain

$$\begin{aligned} \left| \sum_{k=p}^q \lambda_k \Delta v_k \right| &\leq \max_{p \leq r \leq q} \left| \sum_{k=r}^q \Delta v_k \right| \left(\sum_{k=p+1}^q |\lambda_k - \lambda_{k-1}| + |\lambda_p| \right) \\ &= \lambda_q \max_{p \leq r \leq q} \left| \sum_{k=r}^q \Delta v_k \right|. \quad \square \end{aligned}$$

Lemma 5. Let (p_n) satisfy the condition (6). If (u_n) is slowly oscillating, then $(V_{n,p}^{(0)}(\Delta u))$ is slowly oscillating and bounded.

Proof. Let (6) be satisfied. It is easy to see that if (u_n) is slowly oscillating, then $V_{n,p}^{(0)}(\Delta u) = O(1)$ by choosing $\lambda_k = P_{k-1}$ in Lemma 4. By the identity $V_{n,p}^{(0)}(\Delta u) = \frac{p_{n-1}}{p_n} \Delta \sigma_{n,p}^{(1)}(u)$, we have $\Delta \sigma_{n,p}^{(1)}(u) = \frac{p_n}{p_{n-1}} V_{n,p}^{(0)}(\Delta u)$. Then, we have

$$\begin{aligned} |\sigma_{n,p}^{(1)}(u) - \sigma_{n-1,p}^{(1)}(u)| &\leq \frac{p_n}{P_{n-1}} C, \\ |\sigma_{k,p}^{(1)}(u) - \sigma_{n,p}^{(1)}(u)| &\leq \left(\frac{p_k}{P_{k-1}} + \frac{p_{k-1}}{P_{k-2}} + \cdots + \frac{p_{n+1}}{P_n} \right) C \\ |\sigma_{k,p}^{(1)}(u) - \sigma_{n,p}^{(1)}(u)| &\leq \left(\frac{P_k - P_n}{P_n} \right) C \end{aligned}$$

for some $C > 0$. Taking max of both sides of the inequality over k , where k runs from $n+1$ to $[\lambda n]$, we obtain

$$\max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta \sigma_{j,p}^{(1)}(u) \right| \leq \left(\frac{P_{[\lambda n]} - P_n}{P_n} \right) C.$$

Taking lim sup of both sides of the inequality above as $n \rightarrow \infty$, we have

$$\limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta \sigma_{j,p}^{(1)}(u) \right| \leq \limsup_{n \rightarrow \infty} \left(\frac{P_{[\lambda n]} - P_n}{P_n} \right) C. \quad (9)$$

Taking the limit of both sides as $\lambda \rightarrow 1^+$, we obtain by (6)

$$\lim_{\lambda \rightarrow 1^+} \limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta \sigma_{j,p}^{(1)}(u) \right| \leq 0. \quad (10)$$

It follows from (10) that $(\sigma_{n,p}^{(1)}(u))$ is slowly oscillating. Since (u_n) is slowly oscillating, $(V_{n,p}^{(0)}(\Delta u))$ is slowly oscillating by the weighted Kronecker identity. \square

The following difference $u_n - \sigma_{n,p}^{(1)}(u)$ for a sequence $u = (u_n)$ is well known and an easy consequence of (2).

Lemma 6 ([6]). Let (u_n) be a sequence of real numbers. For $\lambda > 1$ and sufficiently large n ,

$$u_n - \sigma_{n,p}^{(1)}(u) = \frac{P_{[\lambda n]}}{P_{[\lambda n]} - P_n} \left(\sigma_{[\lambda n],p}^{(1)}(u) - \sigma_{n,p}^{(1)}(u) \right) - \frac{1}{P_{[\lambda n]} - P_n} \sum_{k=n+1}^{[\lambda n]} p_k (u_k - u_n)$$

where $[\lambda n]$ denotes the integer part of λn .

3. Proof of Theorem 1

By hypothesis, (u_n) is (J, p) summable to s . Since (u_n) is slowly oscillating sequence, we have from Lemma 5 that $V_{n,p}^{(0)}(\Delta u) = O(1)$. We thus obtain $V_{n,p}^{(0)}(\Delta u) \geq -C$ for some $C > 0$. From Theorem 3, (u_n) is (\bar{N}, p) summable to s . It follows from the weighted Kronecker identity that $(V_{n,p}^{(1)}(\Delta u))$ is convergent to zero. Since (u_n) is slowly oscillating sequence, $(V_{n,p}^{(0)}(\Delta u))$ is slowly oscillating by Lemma 5. It suffices to show that $V_{n,p}^{(0)}(\Delta u) = o(1)$. By Lemma 6, we get

$$\begin{aligned} V_{n,p}^{(0)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) &= \frac{P_{[\lambda n]}}{P_{[\lambda n]} - P_n} \left(V_{[\lambda n],p}^{(1)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right) - \frac{1}{P_{[\lambda n]} - P_n} \sum_{k=n+1}^{[\lambda n]} p_k (V_{k,p}^{(0)}(\Delta u) - V_{n,p}^{(0)}(\Delta u)) \\ &= \frac{P_{[\lambda n]}}{P_{[\lambda n]} - P_n} \left(V_{[\lambda n],p}^{(1)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right) - \frac{1}{P_{[\lambda n]} - P_n} \sum_{k=n+1}^{[\lambda n]} p_k \sum_{j=n+1}^k \Delta V_{j,p}^{(0)}(\Delta u). \end{aligned}$$

By the triangle inequality, we have

$$\left| V_{n,p}^{(0)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right| \leq \left| \frac{P_{[\lambda n]}}{P_{[\lambda n]} - P_n} \left(V_{[\lambda n],p}^{(1)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right) \right| + \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta V_{j,p}^{(0)}(\Delta u) \right|. \quad (11)$$

Taking lim sup of both sides of the inequality (11), we obtain

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left| V_{n,p}^{(0)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right| &\leq \limsup_{n \rightarrow \infty} \frac{P_{[\lambda n]}}{P_{[\lambda n]} - P_n} \limsup_{n \rightarrow \infty} \left| V_{[\lambda n],p}^{(1)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right| \\ &\quad + \limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta V_{j,p}^{(0)}(\Delta u) \right|. \end{aligned} \quad (12)$$

Since the first term on the right-hand side of the inequality (12) vanishes by (6), we have,

$$\limsup_{n \rightarrow \infty} \left| V_{n,p}^{(0)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right| \leq \limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta V_{j,p}^{(0)}(\Delta u) \right|.$$

Since $(V_{n,p}^{(0)}(\Delta u))$ is slowly oscillating, taking the limit of both sides as $\lambda \rightarrow 1^+$, we obtain $\limsup_{n \rightarrow \infty} \left| V_{n,p}^{(0)}(\Delta u) - V_{n,p}^{(1)}(\Delta u) \right| \leq 0$, which implies that $V_{n,p}^{(0)}(\Delta u) = o(1)$. Since (u_n) is (\bar{N}, p) summable to s , (u_n) is convergent to s by the weighted Kronecker identity. \square

4. Proof of Theorem 2

By hypothesis, (u_n) is $(J, p)(\bar{N}_m, p)$ summable to s . That is, $(\sigma_{n,p}^{(m)}(u))$ is (J, p) summable to s for some nonnegative integer m . Since (u_n) is slowly oscillating, we have from Lemma 5 that $V_{n,p}^{(0)}(\Delta u) = O(1)$. From here we get $V_{n,p}^{(m)}(\Delta u) = O(1)$. We thus obtain $V_{n,p}^{(m)}(\Delta u) \geq -C$ for some $C > 0$. From Theorem 3, $(\sigma_{n,p}^{(m)}(u))$ is (\bar{N}, p) summable to s . Therefore we have $(\sigma_{n,p}^{(m+1)}(u))$ is convergent to s . Since (u_n) is slowly oscillating sequence, $(\sigma_{n,p}^{(m)}(u))$ is slowly oscillating from Lemma 5. It suffices to show that $(\sigma_{n,p}^{(m)}(u))$ is convergent to s . By Lemma 6, we get

$$\begin{aligned} \sigma_{n,p}^{(m)}(u) - \sigma_{n,p}^{(m+1)}(u) &= \frac{P_{[\lambda n]}}{P_{[\lambda n]} - P_n} \left(\sigma_{[\lambda n],p}^{(m+1)}(u) - \sigma_{n,p}^{(m+1)}(u) \right) - \frac{1}{P_{[\lambda n]} - P_n} \sum_{k=n+1}^{[\lambda n]} p_k (\sigma_{k,p}^{(m)}(u) - \sigma_{n,p}^{(m)}(u)) \\ &= \frac{P_{[\lambda n]}}{P_{[\lambda n]} - P_n} \left(\sigma_{[\lambda n],p}^{(m+1)}(u) - \sigma_{n,p}^{(m+1)}(u) \right) - \frac{1}{P_{[\lambda n]} - P_n} \sum_{k=n+1}^{[\lambda n]} p_k \sum_{j=n+1}^k \Delta \sigma_{j,p}^{(m)}(u). \end{aligned}$$

By the triangle inequality, we have

$$\left| \sigma_{n,p}^{(m)}(u) - \sigma_{n,p}^{(m+1)}(u) \right| \leq \left| \frac{P_{[\lambda n]} }{P_{[\lambda n]} - P_n} \left(\sigma_{[\lambda n],p}^{(m+1)}(u) - \sigma_{n,p}^{(m+1)}(u) \right) \right| + \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta \sigma_{j,p}^{(m)}(u) \right|. \quad (13)$$

Taking \limsup of both sides of the inequality (13), we obtain

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left| \sigma_{n,p}^{(m)}(u) - \sigma_{n,p}^{(m+1)}(u) \right| &\leq \limsup_{n \rightarrow \infty} \frac{P_{[\lambda n]} }{P_{[\lambda n]} - P_n} \limsup_{n \rightarrow \infty} \left| \sigma_{[\lambda n],p}^{(m+1)}(u) - \sigma_{n,p}^{(m+1)}(u) \right| \\ &\quad + \limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta \sigma_{j,p}^{(m)}(u) \right|. \end{aligned} \quad (14)$$

Since the first term on the right-hand side of the inequality (14) vanishes by (6), we have

$$\limsup_{n \rightarrow \infty} \left| \sigma_{n,p}^{(m)}(u) - \sigma_{n,p}^{(m+1)}(u) \right| \leq \limsup_{n \rightarrow \infty} \max_{n+1 \leq k \leq [\lambda n]} \left| \sum_{j=n+1}^k \Delta \sigma_{j,p}^{(m)}(u) \right|.$$

Since $(\sigma_{n,p}^{(m)}(u))$ is slowly oscillating, taking the limit of both sides as $\lambda \rightarrow 1^+$, we obtain $\limsup_{n \rightarrow \infty} \left| \sigma_{n,p}^{(m)}(u) - \sigma_{n,p}^{(m+1)}(u) \right| \leq 0$, which implies that $(\sigma_{n,p}^{(m)}(u))$ is convergent to s . This means that $(\sigma_{n,p}^{(m-1)}(u))$ is (\bar{N}, p) summable to s . Therefore, we have $(\sigma_{n,p}^{(m-1)}(u))$ is (J, p) summable to s . Repeating the above process $m - 1$ times we obtain (u_n) is (J, p) summable to s . This completes the proof by Theorem 1. \square

References

- [1] K. Ishiguro, A tauberian theorem for (J, p_n) summability, Proc. Japan Acad. 40 (1964) 807–812.
- [2] Č.V. Stanojević, Analysis of Divergence: Control and Management of Divergent Process, Graduate Research Seminar Lecture Notes, edited by İ. Çanak, University of Missouri–Rolla, Fall 1998.
- [3] G.H. Hardy, Divergent Series, Clarendon Press, Oxford, 1949.
- [4] İ. Çanak, A short proof of the generalized littlewood tauberian theorem, Appl. Math. Lett. 23 (7) (2010) 818–820.
- [5] G.A. Mikhailin, Theorem of tauberian type for (J, p_n) summation methods, Ukrain. Mat. Zh. 29 (1977) 763–770. English translation: Ukrain. Math. J. 29 (1977), 564–569.
- [6] F. Moricz, B.E. Rhoades, Necessary and sufficient tauberian conditions for certain weighted mean methods of summability, Acta Math. Hungar. 66 (1–2) (1995) 105–111.